

MAP 3306 eMath2 **Quiz 8** 20 Jul 2007 Name: \_\_\_\_\_

**Directions:** Show **ALL** work for credit; Give **EXACT** answers when possible; **SIMPLIFY** your answers;

1. Use Fourier transforms to find the solution  $u(x, t)$  of the PDE  $u_x + u_t + 2u = 0$   $u(x, 0) = f(x)$ . The Equations on the other side of the page might be helpful.

- (1)  $\mathcal{F}[f(x)] = \hat{f}(w)$  or simply  $\mathcal{F}[f] = \hat{f}$
- (2)  $\mathcal{F}^{-1}[\hat{f}(w)] = f(x)$  or simply  $\mathcal{F}^{-1}[\hat{f}] = f$
- (3)  $\mathcal{F}[f(x)](w) = \hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{-iwx} dx$
- (4)  $\mathcal{F}^{-1}[\hat{f}(w)](x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(w)e^{iwx} dw$
- (5)  $\mathcal{F}[u(x, t)](w, t) = \hat{u}(w, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} u(x, t)e^{-iwx} dx$
- (6)  $\mathcal{F}^{-1}[\hat{u}(w, t)](x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{u}(w, t)e^{iwx} dw$
- (7)  $\mathcal{F}[af(x) + bg(x)](w) = a\hat{f}(w) + b\hat{g}(w)$
- (8)  $\mathcal{F}[f'(x)](w) = iw\hat{f}(w)$
- (9)  $\mathcal{F}[f''(x)](w) = -w^2\hat{f}(w)$
- (10)  $\mathcal{F}\left[\frac{\partial}{\partial x}u(x, t)\right](w, t) = iw\hat{u}(w, t)$
- (11)  $\mathcal{F}\left[\frac{\partial^2}{\partial x^2}u(x, t)\right](w, t) = -w^2\hat{u}(w, t)$
- (12)  $\mathcal{F}\left[\frac{\partial}{\partial t}u(x, t)\right](w, t) = \frac{\partial}{\partial t}\hat{u}(w, t)$
- (13)  $\mathcal{F}\left[\frac{\partial^2}{\partial t^2}u(x, t)\right](w, t) = \frac{\partial^2}{\partial t^2}\hat{u}(w, t)$
- (14)  $[f * g](x) = \int_{-\infty}^{\infty} f(w)g(x - w) dw = [g * f](x) = \int_{-\infty}^{\infty} f(x - w)g(w) dw$
- (15)  $\mathcal{F}[f * g] = \sqrt{2\pi}\hat{f}\hat{g}$
- (16)  $f(x - a) = \mathcal{F}^{-1}[e^{-iwa}\hat{f}(w)]$
- (17)  $\mathcal{F}[\exp(-ax^2)] = \frac{1}{\sqrt{2a}} \exp\left(\frac{-w^2}{4a}\right)$
- (18)  $\sin wa = \frac{e^{iwa} - e^{-iwa}}{2i}$
- (19)  $\cos wa = \frac{e^{iwa} + e^{-iwa}}{2}$
- (20)  $\frac{1}{\sqrt{2\pi}} \int_{-a}^a e^{-iwx} dx = \sqrt{\frac{2}{\pi}} \frac{\sin aw}{w}$
- (21)  $\mathcal{F}\left[\frac{\sin ax}{x}\right] = \sqrt{\frac{\pi}{2}}$  if  $|w| < a$ ; 0 otherwise
- (22)  $\frac{1}{\sqrt{2\pi}} \int_0^{\infty} e^{-ax} e^{-iwx} dx = \frac{1}{\sqrt{2\pi}(a + iw)}$