

# Complex Homework Summer 2014

Based on Brown and Churchill 7th Edition

June 21, 2014

## Contents

1	hw1, Complex Arithmetic, Conjugates, Polar Form	2
2	hw2 nth roots, Domains, Functions	2
3	hw3 Images, Transformations	3
4	hw4 Limits	3
5	hw5 Unbounded	4
6	hw6 Derivatives, Cauchy-Riemann	4
7	hw7 Exp and Log	5
8	hw8 Log and log	5
9	hw9 Principal values, Integrals over a Real Variable	6
10	hw10 Contour Integrals	6
11	hw11 More on Contour Integrals	7
12	hw12 Path independence	7
13	hw13 Cauchy Goursat	8
14	hw14 Applications of Cauchy Integral Formula	8
15	hw15 Liouville	9
16	hw16 Series	9
17	hw17 Taylor Series	10
18	hw18 Laurent Series	10
19	hw19 Derivative of Series, Substituting, Poles, Residues	11
20	hw20 Singular points	12
21	hw21 Residues, Poles, Order of a Pole	12

**22 hw22 Computing Integrals** **13**

**23 hw23 Poles and Zeros** **13**

**24 hw24 Cool Integrals** **13**

These are problems will be due both daily and at the end of classes. This PDF file was created on June 21, 2014.

## 1 hw1, Complex Arithmetic, Conjugates, Polar Form

1. (BC3.1) Reduce each of these 3 expressions to a real number

$$\frac{1+2i}{3-4i} + \frac{2-i}{5i} \quad \frac{5i}{(1-i)(2-i)(3-i)} \quad \text{and} \quad (1-i)^4$$

2. (BC4.1) In each case locate  $z_1 + z_2$  and  $z_1 - z_2$  vectorially

$$\begin{array}{ll} z_1 = 2i, z_2 = \frac{2}{3} - i & z_1 = (-\sqrt{3}, 0), z_2 = (\sqrt{3}, 0) \\ z_1 = (-3, 1), z_2 = (1, 4) & z_1 = x_1 + iy_1, z_2 = x_1 - iy_1 \end{array}$$

3. (BC4.4) Sketch the set of points determined by each equation

$$|z - 1 + i| = 1 \quad |z + i| \leq 3 \quad \text{and} \quad |z + 4i| \geq 4$$

4. (BC5.3,4) Verify  $\overline{z_1 - z_2} = \overline{z_1} - \overline{z_2}$ ,  $\overline{z_1 z_2} = \overline{z_1} \overline{z_2}$ ,  $\overline{z_1 z_2 z_3} = \overline{z_1} \overline{z_2} \overline{z_3}$  and  $\overline{z^4} = \overline{z}^4$ .

5. (BC5.5) Verify

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad (z_2 \neq 0)$$

6. (BC5.15) Show that the hyperbola  $x^2 - y^2 = 1$  can be written  $z^2 + \overline{z}^2 = 2$

7. (BC7.1) Find the principal argument  $\text{Arg } z$  for both

$$z = \frac{i}{-2 - 2i} \quad \text{and} \quad z = (\sqrt{3} - i)^6$$

8. (BC7.2) Show  $|e^{i\theta}| = 1$  and  $\overline{e^{i\theta}} = e^{-i\theta}$

9. (BC7.15) Use de Moivre's formula to derive the following trig identities.

$$\cos 3\theta = \cos^3 \theta - 3 \cos \theta \sin^2 \theta = 4 \cos^3 \theta - 3 \cos \theta$$

$$\sin 3\theta = 3 \cos^2 \theta \sin \theta - \sin^3 \theta = 3 \sin \theta - 4 \sin^3 \theta$$

## 2 hw2 nth roots, Domains, Functions

1. (BC7.7) Show if  $\Re z_1 > 0$  and  $\Re z_2 > 0$  then  $\text{Arg}(z_1 z_2) = \text{Arg } z_1 + \text{Arg } z_2$

2. (BC9.1) Find the square roots of  $2i$  and  $1 - i\sqrt{3}$  expressed in rectangular form

3. (BC9.3) Find all of the roots in rectangle coordinates of  $(-1)^{1/3}$  and  $8^{1/6}$ .

4. (BC9.6) Find the 4 roots of  $p(z) = z^4 + 4 = 0$  and use them to factor  $p(z)$  into quadratic factors with real coefficients.
5. (BC10.1-3) Sketch the 6 sets and determine which are domains, which are bounded, which are neither open nor closed:
- $$\begin{array}{lll} |z - 2 + i| \leq 1 & |2z + 3| > 4 & \Im z > 1 \\ \Im z = 1 & 0 \leq \arg z \leq \pi/4 (z \neq 0) & |z - 4| \leq |z| \end{array}$$

6. (BC10.4) Find the closure of the 4 sets:

$$-\pi < \arg z < \pi (z \neq 0) \quad |\Re z| < |z| \quad \Re\left(\frac{1}{z}\right) \leq \frac{1}{2} \quad \text{and} \quad \Re(z^2) > 0$$

7. (BC11.1) For each function, describe the domain that is understood:

$$f(z) = \frac{1}{z^2 + 1} \quad f(z) = \text{Arg}\left(\frac{1}{z}\right) \quad f(z) = \frac{z}{z + \bar{z}} \quad \text{and} \quad f(z) = \frac{1}{1 - |z|^2}$$

8. (BC11.2) Write  $z^3 + z + 1$  as  $u(x, y) + iv(x, y)$
9. (BC11.3) Write and simplify  $f(z) = x^2 - y^2 - 2y + i(2x - 2xy)$  in terms of  $z$  using  $x = (z + \bar{z})/2$  and  $y = (z - \bar{z})/2i$
10. (BC11.4) Write  $f(z) = z + 1/z (z \neq 0)$  in the form  $u(r, \theta) + iv(r, \theta)$

### 3 hw3 Images, Transformations

1. (BC13.1) Find a domain in the  $z$ -plane whose image under the transformation  $w = z^2$  is the square domain in the  $w$ -plane bounded by the lines  $u = 1, u = 2, v = 1, v = 2$
2. (BC13.3) Sketch the region onto which the sector  $r \leq 1, 0 \leq \theta \leq \pi/4$  is mapped by the 3 transformations  $w = z^2, w = z^3$ , and  $w = z^4$
3. (BC13.4) Show that lines  $ay = x (a \neq 0)$  are mapped onto the spirals  $\rho = \exp(a\theta)$  under the transformation  $w = \exp z$ , where  $w = \rho \exp(i\phi)$
4. (BC13.7) Find the image of the semi-infinite strip  $x \geq 0, 0 \leq y \leq \pi$  under the transformation  $w = \exp z$ . Label the corresponding portions of the boundaries.
5. (BC13.8) Graphically indicate the vector fields represented by  $w = iz$  and  $w = z/|z|$

### 4 hw4 Limits

1. (BC17.3) Find the limits.  $n$  is a positive integer,  $P(z)$  and  $Q(z)$  are polynomials with  $Q(z_0) \neq 0$

$$\lim_{z \rightarrow z_0} \frac{1}{z^n} (z_0 \neq 0) \quad \lim_{z \rightarrow i} \frac{iz^3 - 1}{z + i} \quad \text{and} \quad \lim_{z \rightarrow z_0} \frac{P(z)}{Q(z)}$$

2. (BC17.5) Show that the following limit does not exist

$$\lim_{z \rightarrow 0} \left(\frac{z}{\bar{z}}\right)^2$$

3. (BC17.10) Use a theorem to show:

$$\lim_{z \rightarrow \infty} \frac{4z^2}{(z-1)^2} = 4 \quad \lim_{z \rightarrow 1} \frac{1}{(z-1)^3} = \infty \quad \text{and} \quad \lim_{z \rightarrow \infty} \frac{z^2+1}{z-1} = \infty$$

4. (BC17.11) Suppose  $ad - bc \neq 0$  and let:

$$T(z) = \frac{az+b}{cz+d}$$

Use a theorem to show

$$\lim_{z \rightarrow \infty} T(z) = \infty \text{ (if } c = 0) \quad \lim_{z \rightarrow \infty} T(z) = \frac{a}{c} \text{ (if } c \neq 0) \quad \text{and} \quad \lim_{z \rightarrow -d/c} T(z) = \infty \text{ (if } c \neq 0)$$

## 5 hw5 Unbounded

1. (BC17.13) (Show that a set  $S$  is unbounded if and only if every neighborhood of the point at infinity contains at least one point of  $S$ .)

## 6 hw6 Derivatives, Cauchy-Riemann

1. (BC19.1) Find  $f'(z)$  when

$$f(z) = 3z^2 - 2z + 4 \quad f(z) = (1 - 4z^2)^3 \quad f(z) = \frac{z-1}{2z+1} \quad (z \neq -\frac{1}{2}) \quad \text{and} \quad f(z) = \frac{(1+z^2)^4}{z^2} \quad (z \neq 0)$$

2. (BC19.2) Show if  $P(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$  then  $P'(z) = a_1 + 2a_2z + \cdots + na_nz^{n-1}$  and hence

$$a_0 = P(0), \quad a_1 = \frac{P'(0)}{1!}, \quad a_2 = \frac{P''(0)}{2!}, \quad \dots \quad a_n = \frac{P^{(n)}(0)}{n!}$$

3. (BC19.9) Let  $f$  denote the function whose values are

$$f(z) = \begin{cases} \bar{z}^2/z & \text{when } z \neq 0 \\ 0 & \text{when } z = 0 \end{cases}$$

Show that if  $z = 0$ , then  $\Delta w/\Delta z = 1$  at each nonzero point on the real and imaginary axes in the  $\Delta z$  or  $\Delta x\Delta y$ -plane. Then show then  $\Delta w/\Delta z = -1$  at each nonzero point along the line  $y = x$ . Conclude that  $f'(0)$  does not exist.

4. (BC22.6) Let  $f$  denote the function above. Show that the Cauchy-Riemann equations are satisfied at the origin  $z = (0, 0)$
5. (BC22.1) Use a theorem to show that  $f'(z)$  does not exist at any point for each function:

$$f(z) = \bar{z} \quad f(z) = z - \bar{z} \quad f(z) = 2x + ixy^2 \quad \text{and} \quad f(x) = e^x e^{-iy}$$

6. (BC22.2) Use a theorem to show that  $f'(z)$  and its derivative  $f''(z)$  exist everywhere and find  $f''(z)$ .

$$f(z) = iz + 2 \quad f(z) = e^{-x} e^{-iy} \quad f(z) = z^3 \quad \text{and} \quad f(z) = \cos x \cosh y - i \sin x \sinh y$$

7. Extra Credit (BC22.10) Recall  $z = x + iy$  implies  $x = (z + \bar{z})/2$  and  $y = (z - \bar{z})/2i$ . Use the formal chain rule to show

$$\frac{\partial F}{\partial \bar{z}} = \frac{\partial F}{\partial x} \frac{\partial x}{\partial \bar{z}} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial F}{\partial x} + i \frac{\partial F}{\partial y} \right)$$

Define the operator

$$\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$$

and apply it to  $u(x, y) + iv(x, y)$  to obtain the complex form of the Cauchy-Reimann equations  $\partial f / \partial \bar{z} = 0$ .

## 7 hw7 Exp and Log

- (BC28.1) Show that  $\exp(2 \pm 3\pi i) = -e^2$ ,  $\exp((2 + \pi i)/4) = (1 + i)\sqrt{e/2}$  and  $\exp(z + \pi i) = -\exp z$ .
- (BC28.2) State why the function  $2z^2 - 3 - ze^z + e^{-z}$  is entire.
- (BC28.3) Show  $f(z) = \exp \bar{z}$  is not analytic anywhere.
- (BC28.7) Prove  $|\exp(-2z)| < 1$  if and only if  $\Re z > 0$ .
- (BC28.8) Find all values of  $z$  such that  $e^z = -2$ , or  $e^z = 1 + \sqrt{3}i$  or  $\exp(2z - 1) = 1$
- (BC28.10) Show that if  $e^z$  is real, then  $\Im z = n\pi$  ( $n = 0, \pm 1, \pm 2, \dots$ ). If  $e^z$  is pure imaginary, what restriction is placed on  $z$ ?
- (BC30.1) Show that  $\text{Log}(-ei) = 1 - \frac{\pi}{2}i$  and  $\text{Log}(1 - i) = \frac{1}{2} \ln 2 - \frac{\pi}{4}i$ .

## 8 hw8 Log and log

- (BC30.2) Verify for  $n = 0, \pm 1, \pm 2, \dots$ :

$$\log e = 1 + 2n\pi i \quad \log i = \left(2n + \frac{1}{2}\right)\pi i \quad \text{and} \quad \log(-1 + \sqrt{3}i) = \ln 2 + 2\left(n + \frac{1}{3}\right)\pi i$$

- (BC30.3) Show that  $\text{Log}(1 + i)^2 = 2\text{Log}(1 + i)$  and  $\text{Log}(-1 + i)^2 \neq 2\text{Log}(-1 + i)$ .
- (BC30.5) Show that the set of values of  $\log(i^{1/2})$  is  $\{(n + \frac{1}{4})\pi i : n = 0, \pm 1, \pm 2, \dots\}$  and that the same is true of  $(1/2)\log i$ .
- (BC30.6) Given that the branch  $\log z = \ln r + i\theta$  ( $r > 0, \alpha < \theta < \alpha + 2\pi$ ) of the logarithmic function is analytic at each point  $z$  in the stated domain, obtain its derivative by differentiating each side of the identity  $\exp(\log z) = z$  and using the chain rule.
- (BC30.7) Find all the roots of the equation  $\log z = i\pi/2$ .
- (BC30.9) Show that  $\text{Log}(z - i)$  is analytic everywhere except on the half line  $y = 1$  ( $x \leq 0$ ). Show

$$\frac{\text{Log}(z + 4)}{z^2 + i}$$

is analytic everywhere except at the points  $\pm(1 - i)/\sqrt{2}$  and on the portion  $x \leq -4$  of the real axis.

## 9 hw9 Principal values, Integrals over a Real Variable

- (BC31.1) Show if  $\Re z_1 > 0$  and  $\Re z_2 > 0$  then  $\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2$ .
- (BC31.2) Show that for any two complex numbers  $z_1$  and  $z_2$ ,  $\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2 + 2N\pi i$  where  $N$  has one of the values  $0, \pm 1$ .
- (BC32.1) Show that when  $n = 0, \pm 1, \pm 2, \dots$

$$(1+i)^i = \exp\left(-\frac{\pi}{4} + 2n\pi\right) \exp\left(\frac{i}{2} \ln 2\right) \quad \text{and} \quad (-1)^{1/\pi} = e^{(2n+1)i}$$

- (BC32.2) Find the principal values of each expression:

$$i^i \quad \left[\frac{e}{2}(-1 - \sqrt{3}i)\right]^{3\pi i} \quad \text{and} \quad (1-i)^{4i}$$

- (BC32.5) Show that the principal  $n$ -th root of a nonzero complex number  $z_0$  is the same as the principal value of  $z_0^{1/n}$  that was previously defined.
- (BC32.8) Let  $c, d, z$  be complex numbers with  $z \neq 0$ . Prove that if all the powers involved are principal values, then

$$\frac{1}{z^c} = z^{-c} \quad (z^c)^n = z^{cn} \quad (n = 1, 2, \dots) \quad z^c z^d = z^{c+d} \quad \text{and} \quad \frac{z^c}{z^d} = z^{c-d}$$

- (BC37.2) Evaluate

$$\int_1^2 \left(\frac{1}{t} - i\right)^2 dt \quad \int_0^{\pi/6} e^{i2t} dt \quad \text{and} \quad \int_0^\infty e^{-zt} dt \quad (\Re z > 0)$$

- (BC37.5) Let  $w(t)$  be a continuous complex-valued function of  $t$  defined on an interval  $a \leq t \leq b$ . By considering the special case  $w(t) = e^{it}$  on the interval  $0 \leq t \leq 2\pi$ , show that it is not always true that there is a number  $c$  in the interval  $a < t < b$  such that

$$\int_a^b w(t) dt = w(c)(b-a)$$

## 10 hw10 Contour Integrals

- (BC38.2) Let  $C$  denote the right-hand half of the circle  $|z| = 2$ , in the counterclockwise direction and note that two parametric representations for  $C$  are

$$z = z(\theta) = 2e^{i\theta} \quad \left(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}\right)$$

and

$$z = Z(y) = \sqrt{4-y^2} + iy \quad (-2 \leq y \leq 2)$$

Verify that  $Z(y) = z[\phi(y)]$ , where

$$\phi(y) = \arctan \frac{y}{\sqrt{4-y^2}} \quad \left(-\frac{\pi}{2} \leq \arctan t \leq \frac{\pi}{2}\right)$$

Also, show that this function  $\phi$  has a positive derivative, as required in the conditions following (9) Sec 38.

2. (BC40.1,2,3,5,6) Evaluate

$$\int_C f(z) dz$$

for the given  $f(z)$  and contour  $C$

- $f(z) = (z + 2)/z$      $C$  is  $z = 2e^{i\theta}$  ( $0 \leq \theta \leq \pi$ )  
 $f(z) = (z + 2)/z$      $C$  is  $z = 2e^{i\theta}$  ( $\pi \leq \theta \leq 2\pi$ )  
 $f(z) = (z + 2)/z$      $C$  is  $z = 2e^{i\theta}$  ( $0 \leq \theta \leq 2\pi$ )  
 $f(z) = z + 1$          $C$  is  $z = 1 + e^{i\theta}$  ( $\pi \leq \theta \leq 2\pi$ )  
 $f(z) = z + 1$          $C$  is  $z = t$  ( $0 \leq t \leq 2$ )  
 $f(z) = \pi \exp(\pi \bar{z})$      $C$  is square from  $0, 1, 1 + i, i$   
 $f(z) = 1$              $C$  is arbitrary curve from  $z_1$  to  $z_2$   
 $f(z) = z^{-1+i}$          $C$  is  $|z| = 1$  positively oriented  
                           use branch  $\exp[(-1 + i) \log z]$  ( $|z| > 0, 0 < \arg z < 2\pi$ )

3. (BC40.10) Let  $C_0$  denote the circle  $|z - z_0| = R$  taken counterclockwise. Use the parametric representation  $z = z_0 + Re^{i\theta}$  ( $-\pi \leq \theta \leq \pi$ ) for  $C_0$  to derive the following integration formula's:

$$\int_{C_0} \frac{dz}{z - z_0} = 2\pi i \quad \text{and} \quad \int_{C_0} (z - z_0)^{n-1} dz = 0 \quad (n = \pm 1, \pm 2, \dots)$$

## 11 hw11 More on Contour Integrals

1. (BC41.4) Let  $C_R$  denote the upper half of the circle  $|z| = R$  ( $R > 2$ ), taken in the counterclockwise direction. Show that

$$\left| \int_{C_R} \frac{2z^2 - 1}{z^4 + 5z^2 + 4} dz \right| \leq \frac{\pi R(2R^2 + 1)}{(R^2 - 1)(R^2 - 4)}$$

2. (BC43.1) Use an antiderivative to show that, for every contour  $C$  extending from a point  $z_1$  to a point  $z_2$ ,

$$\int_C z^n dz = \frac{1}{n+1} (z_2^{n+1} - z_1^{n+1}) \quad (n = 0, 1, \dots)$$

3. (BC43.2) By finding an antiderivative, evaluate each of these integrals, where the path is any contour between the indicated limits of integration.

$$\int_i^{i/2} e^{\pi z} dz \quad \int_0^{\pi+2i} \cos\left(\frac{z}{2}\right) dz \quad \text{and} \quad \int_1^3 (z-2)^3 dz$$

## 12 hw12 Path independence

1. (BC43.3) Use a theorem to show

$$\int_{C_0} (z - z_0)^{n-1} dz = 0 \quad (n = \pm 1, \pm 2, \dots)$$

when  $C_0$  is any closed contour which does not pass through the point  $z_0$ .

2. (BC43.4) Let  $C_1$ , (resp.  $C_2$ ), be any contour from  $z = -3$  to  $z = 3$  that except for its end points, lies above (resp. below) the  $x$ -axis. Find an antiderivative  $F_2(z)$  of the branch  $f_2(z)$  of

$$z^{1/2} = \sqrt{r}e^{i\theta/2} \quad (r > 0, \frac{\pi}{2} < \theta < \frac{5\pi}{2})$$

to show that the integral

$$\int_{C_2} z^{1/2} dz$$

has value  $2\sqrt{3}(-1 + i)$ . Note that the value of the integral of the function

$$z^{1/2} = \sqrt{r}e^{i\theta/2}$$

around the closed contour  $C_2 - C_1$  in that example is, therefore  $-4\sqrt{3}$  given that

$$\int_{C_1} z^{1/2} dz = 2\sqrt{3}(1 + i)$$

. (Lots of parts from example 43.4.)

### 13 hw13 Cauchy Goursat

- (BC46.1) Apply the Cauchy-Goursat theorem to show that

$$\int_C f(z) dz = 0$$

when the contour  $C$  is the circle  $|z| = 1$ , in either direction and when

$$\begin{array}{lll} f(z) = \frac{z^2}{z-3} & f(z) = ze^{-z} & f(z) = \frac{1}{z^2 + 2z + 2} \\ f(z) = \operatorname{sech} z & f(z) = \tan z & f(z) = \operatorname{Log}(z + 2) \end{array}$$

- (BC46.2) Let  $C_1$  be the positively oriented circle  $|z| = 4$  and let  $C_2$  be the positively oriented boundary of the square whose sides lie along the lines  $x = \pm 1, y = \pm 1$ . Point out why

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

when

$$f(z) = \frac{1}{3z^2 + 1} \quad f(z) = \frac{z + 2}{\sin(z/2)} \quad \text{and} \quad f(z) = \frac{z}{1 - e^z}$$

- (BC46.3) If  $C$  is the boundary of the rectangle  $0 \leq x \leq 3, 0 \leq y \leq 2$ , described in the positive sense, then

$$\int_C (z - 2 - i)^{n-1} dz = 2\pi i \text{ when } n = 0 \text{ and } 0 \text{ when } n = \pm 1, \pm 2, \dots$$

- (BC46.4) Extra Credit ????

### 14 hw14 Applications of Cauchy Integral Formula

- (BC48.1abc) Let  $C$  denote the positively oriented boundary of the square whose sides lie along the lines  $x = \pm 2, y = \pm 2$ . Evaluate the integrals

$$\int_C \frac{e^{-z} dz}{z - (\pi i/2)} \quad \int_C \frac{\cos z dz}{z(z^2 + 8)} \quad \text{and} \quad \int_C \frac{z dz}{2z + 1}$$



- (BC48.2) Find the integral of  $g(z)$  around the circle  $|z - i| = 2$  in the positive sense when  $g(z) = 1/(z^2 + 4)$  and when  $g(z) = 1/(z^2 + 4)^2$ .
- (BC48.3) Let  $C$  be the circle  $|z| = 3$  described in the positive sense. Show that if

$$g(w) = \int_C \frac{2z^2 - z - 2}{z - w} dz \quad (|w| \neq 3)$$

then  $g(2) = 8\pi i$ . What is the value of  $g(w)$  when  $|w| > 3$ ?

- (BC48.7) Let  $C$  be the unit circle  $z = e^{i\theta}$  ( $-\pi \leq \theta \leq \pi$ ). First show that for any real constant  $a$ ,

$$\int_C \frac{e^{az}}{z} dz = 2\pi i$$

Then write this integral in terms of  $\theta$  to derive the integration formula

$$\int_0^\pi e^{a \cos \theta} \cos(a \sin \theta) d\theta = \pi$$

- (BC48.6) Extra Credit ??? Let  $f$  denote a function that is *continuous* on a simple closed contour  $C$ . Prove the function

$$g(z) = \frac{1}{2\pi i} \int_C \frac{f(\xi) d\xi}{\xi - z}$$

is analytic at each point  $z$  interior to  $C$  and that

$$g'(z) = \frac{1}{2\pi i} \int_C \frac{f(\xi) d\xi}{(\xi - z)^2}$$

at such a point.

## 15 hw15 Liouville

- (BC50.1) Let  $f$  be an entire function such that  $|f(z)| \leq A|z|$  for all  $z$ , where  $A$  is a fixed positive number. Show that  $f(z) = a_1 z$ , where  $a_1$  is a complex constant. [Hint: use Cauchy's inequality to show  $f''(z)$  is zero.]
- (BC50.1) Suppose  $f(z)$  is entire and that the harmonic function  $u(x, y) = \Re f(z)$  has an upper bound  $u_0$ : that is,  $u(x, y) \leq u_0$  for all points  $(x, y)$  in the  $xy$ -plane. Show that  $u(x, y)$  must be constant throughout the plane. [Hint: use Liouville's theorem on  $\exp(f(z))$ .]
- (BC50.4,5) Let a function  $f$  be continuous in a closed bounded region  $R$ , and let it be analytic and not constant throughout the interior of  $R$ . Assuming  $f(z) \neq 0$  anywhere in  $R$ , prove that  $|f(z)|$  has a *minimum value*  $m$  in  $R$  which occurs on the boundary of  $R$  and never in the interior. [Hint: look at  $1/f(z)$ .]  
Use the function  $f(z) = z$  to show that the condition  $f(z) \neq 0$  anywhere is necessary for this conclusion.

## 16 hw16 Series

- (BC52.6) Show if  $\sum_{n=1}^\infty z_n = S$ , then  $\sum_{n=1}^\infty \bar{z}_n = \bar{S}$ .
- (BC52.7) Show for any complex number  $c$  Show if  $\sum_{n=1}^\infty z_n = S$ , then  $\sum_{n=1}^\infty cz_n = cS$ .
- (BC52.8) Show if  $\sum_{n=1}^\infty z_n = S$  and  $\sum_{n=1}^\infty w_n = T$ , then  $\sum_{n=1}^\infty (z_n + w_n) = S + T$ .

## 17 hw17 Taylor Series

1. (BC54.2) Obtain the Taylor

$$e^z = e \sum_{n=0}^{\infty} \frac{(z-1)^n}{n!} \quad (|z-1| < \infty)$$

two ways. First using  $f^{(n)}(1)$  and second by using  $e^z = ee^{z-1}$ .

2. (BC54.3) Find the Maclaurin series expansion for the function

$$f(z) = \frac{z}{z^4 + 9} = \frac{z}{9} \cdot \frac{1}{1 + z^4/9}$$

3. (BC54.5) Derive the Maclaurin series for  $\cos z$  by showing  $f^{(2n)}(0) = (-1)^n$  and  $f^{(2n+1)}(0) = 0$  and by using  $\cos z = (e^{iz} + e^{-iz})/2$ .

4. (BC54.11) Show when  $z \neq 0$ ,

$$\begin{aligned} \frac{e^z}{z^2} &= \frac{1}{z^2} + \frac{1}{z} + \frac{1}{2!} + \frac{z}{3!} + \frac{z^2}{4!} + \cdots \\ \frac{\sin(z^2)}{z^4} &= \frac{1}{z^2} - \frac{z^2}{3!} + \frac{z^6}{5!} - \frac{z^{10}}{7!} + \cdots \end{aligned}$$

5. (BC54.13) Show that when  $0 < |z| < 4$ ,

$$\frac{1}{4z - z^2} = \frac{1}{4z} + \sum_{n=0}^{\infty} \frac{z^n}{4^{n+2}}$$

## 18 hw18 Laurent Series

1. (BC56.1) Find the Laurent series that represents the function  $f(z) = z^2 \sin(1/z^2)$  in the domain  $0 < z < \infty$ .

2. (BC56.2) Derive the Laurent series representation

$$\frac{e^z}{(z+1)^2} = \frac{1}{e} \left[ \sum_{n=0}^{\infty} \frac{(z+1)^n}{(n+2)!} + \frac{1}{z+1} + \frac{1}{(z+1)^2} \right]$$

3. (BC56.3) Find a representation for the function

$$f(z) = \frac{1}{1+z} = \frac{1}{z} \cdot \frac{1}{1+(1/z)}$$

in negative powers of  $z$  that is valid for  $1 < |z| < \infty$ .

4. (BC56.4) Give two Laurent series expansions in powers of  $z$  for the function  $f(z) = 1/[z^2(1-z)]$  and specify the regions in which the expansions are valid. [Hint: about 0 and  $\infty$ ]

5. (BC56.5) Represent the function

$$f(z) = \frac{z+1}{z-1}$$

by both its Maclaurin series (stating where it is valid) and by a Laurent series in the domain  $1 < |z| < \infty$

6. (BC56.6) Show that when  $0 < |z-1| < 2$ ,

$$\frac{z}{(z-1)(z-3)} = -3 \sum_{n=0}^{\infty} \frac{(z-1)^n}{2^{n+2}} - \frac{1}{2(z-1)}$$

## 19 hw19 Derivative of Series, Substituting, Poles, Residues

1. (BC60.1) By differentiating the Maclaurin series representation

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n \quad (|z| < 1)$$

obtain the expressions

$$\frac{1}{(1-z)^2} = \sum_{n=0}^{\infty} (n+1)z^n \quad (|z| < 1)$$

and

$$\frac{2}{(1-z)^3} = \sum_{n=0}^{\infty} (n+1)(n+2)z^n \quad (|z| < 1)$$

2. (BC60.2) By substituting  $1/(1-z)$  for  $z$  in the expansion

$$\frac{1}{(1-z)^2} = \sum_{n=0}^{\infty} (n+1)z^n \quad (|z| < 1)$$

found above, derive the Laurent series representation

$$\frac{1}{z^2} = \sum_{n=2}^{\infty} \frac{(-1)^n(n-1)}{(z-1)^n} \quad (1 < |z-1| < \infty)$$

3. (BC60.3) Find the Taylor series for the function

$$\frac{1}{z} = \frac{1}{2+(z-2)} = \frac{1}{2} \cdot \frac{1}{1+(z-2)/2}$$

about the point  $z_0 = 2$ . Then by differentiating that series term by term, show that

$$\frac{1}{z^2} = \frac{1}{4} \sum_{n=0}^{\infty} (-1)^n (n+1) \left(\frac{z-2}{2}\right)^n \quad (|z-2| < 2)$$

4. (BC61.1) Use multiplication of series to show that

$$\frac{e^z}{z(z^2+1)} = \frac{1}{z} + 1 - \frac{1}{2}z - \frac{5}{6}z^2 + \dots \quad (0 < |z| < 1)$$

5. (BC61.3) Use division to obtain the Laurent series representation

$$\frac{1}{e^z - 1} = \frac{1}{z} - \frac{1}{2} + \frac{1}{12}z - \frac{1}{720}z^3 + \dots \quad (0 < |z| < 2\pi)$$

6. (BC64.1) Find the residue at  $z = 0$  of the functions

$$\frac{1}{z+z^2} \quad z \cos\left(\frac{1}{z}\right) \quad \frac{z - \sin z}{z} \quad \frac{\cot z}{z^4} \quad \text{and} \quad \frac{\sinh z}{z^4(1-z^2)}$$

7. (BC64.2) Use Cauchy's residue theorem to evaluate the integral of each of these functions around the circle  $|z| = 3$  in the positive sense:

$$\frac{\exp(-z)}{z^2} \quad \frac{\exp(-z)}{(z-1)^2} \quad z^2 \exp\left(\frac{1}{z}\right) \quad \text{and} \quad \frac{z+1}{z^2-2z}$$

8. (BC64.3) Use a theorem involving a single residue to evaluate the integral of each of these functions around the circle  $|z| = 2$  in the positive sense.

$$\frac{z^5}{1-z^3} \quad \frac{1}{1+z^2} \quad \text{and} \quad \frac{1}{z}$$

## 20 hw20 Singular points

1. (BC65.1) In each case, write the principal part of the function at its isolated singular point and determine whether that point is a pole, a removable singular point or an essential singular point.

$$z \exp\left(\frac{1}{z}\right) \quad \frac{z^2}{1+z} \quad \frac{\sin z}{z} \quad \frac{\cos z}{z} \quad \text{and} \quad \frac{1}{(2-z)^3}$$

2. (BC65.2) Show that the singular point of each of the following functions is a pole. Determine the order  $m$  of the pole and the corresponding residue  $B$ .

$$\frac{1 - \cosh z}{z^3} \quad \frac{1 - \exp(2z)}{z^4} \quad \text{and} \quad \frac{\exp(2z)}{(z-1)^2}$$

3. (BC65.3) Suppose  $f$  is analytic at  $z_0$  and write  $g(z) = f(z)/(z - z_0)$ . Show that:

- (a) If  $f(z_0) \neq 0$ , then  $z_0$  is a simple pole of  $g$ , with residue  $f(z_0)$ .  
 (b) If  $f(z_0) = 0$ , then  $z_0$  is a removable singular point of  $g$ .

## 21 hw21 Residues, Poles, Order of a Pole

1. (BC65.4) Write the function

$$f(z) = \frac{8a^3 z^2}{(z^2 + a^2)^3} \quad (a > 0)$$

as

$$f(z) = \frac{\phi(z)}{(z - ai)^3} \quad \text{where} \quad \phi(z) = \frac{8a^3 z^2}{(z + ai)^3}$$

Point out why  $\phi(z)$  has a Taylor series representation about  $z = ai$ , and then use it to show that the principal part of  $f$  at that point is

$$\frac{\phi''(ai)/2}{z - ai} + \frac{\phi'(ai)}{(z - ai)^2} + \frac{\phi(ai)}{(z - ai)^3} = -\frac{i/2}{z - ai} - \frac{a/2}{(z - ai)^2} - \frac{a^2 i}{(z - ai)^3}$$

2. (BC67.1) In each case, show that any singular point of the function is a pole. Determine the order  $m$  of the pole and find the corresponding residue  $B$

$$\frac{z^2 + 2}{z - 1} \quad \left(\frac{z}{2z + 1}\right)^3 \quad \text{and} \quad \frac{\exp z}{z^2 + \pi^2}$$

3. (BC67.2) Show that

$$\operatorname{Res}_{z=-1} \frac{z^{1/4}}{z+1} = \frac{1+i}{\sqrt{2}} \quad (|z| > 0, 0 < \arg z < 2\pi)$$

$$\operatorname{Res}_{z=i} \frac{\operatorname{Log} z}{(z^2 + 1)^2} = \frac{\pi + 2i}{8}$$

$$\operatorname{Res}_{z=i} \frac{z^{1/2}}{(z^2 + 1)^2} = \frac{1-i}{8\sqrt{2}} \quad (|z| > 0, 0 < \arg z < 2\pi)$$

4. (BC67.3) Find the value of the integral

$$\int_C \frac{3z^3 + 2}{(z-1)(z^2+9)} dz$$

taken counterclockwise around both circles  $|z-2|=2$  and  $|z|=4$

## 22 hw22 Computing Integrals

1. (BC67.4) Find the value of the integral

$$\int_C \frac{dz}{z^3(z+4)}$$

taken counterclockwise around both circles  $|z| = 2$  and  $|z+2| = 3$

2. (BC69.1) Show that the point  $z = 0$  is a simple pole of the function  $f(z) = \csc z = 1/\sin z$  by a theorem and by computing the Laurent series.

3. (BC69.3a) Show that

$$\operatorname{Res}_{z=z_n} (z \sec z) = (-1)^{n+1} z_n, \text{ where } z_n = \frac{\pi}{2} + n\pi \quad (n = 0, \pm 1, \pm 2, \dots)$$

4. (BC69.4a) Let  $C$  denote the positively oriented circle  $|z| = 2$  and evaluate the integral

$$\int_C \tan z \, dz$$

5. (BC69.5) Let  $C_N$  denote the positive oriented boundary of the square whose edges lie along the lines

$$x = \pm(N + \frac{1}{2})\pi \text{ and } y = \pm(N + \frac{1}{2})\pi$$

where  $N$  is a positive integer. Show that

$$\int_{C_N} \frac{dz}{z^2 \sin z} = 2\pi i \left[ \frac{1}{6} + 2 \sum_{n=1}^N \frac{(-1)^n}{n^2 \pi^2} \right]$$

then using the fact that the value of this integral tends to zero as  $N$  tends to infinity, point out how it follows that

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = \frac{\pi^2}{12}$$

## 23 hw23 Poles and Zeros

1. (BC69.9) Let  $p$  and  $q$  denote functions that are analytic at a point  $z_0$  where  $p(z_0) \neq 0$  and  $q(z_0) = 0$ . Show that if the quotient  $p(z)/q(z)$  has a pole of order  $m$  at  $z_0$ , then  $z_0$  is a zero of order  $m$  of  $q$ .

## 24 hw24 Cool Integrals

1. (BC72.1,2,4) Use residues to evaluate the following integrals

$$\int_0^{\infty} \frac{dx}{x^2+1} \quad \int_0^{\infty} \frac{dx}{(x^2+1)^2} \quad \text{and} \quad \int_0^{\infty} \frac{x^2 dx}{(x^2+1)(x^2+4)}$$

2. (BC74.1,2) Use residues to evaluate the following integrals

$$\int_{-\infty}^{\infty} \frac{\cos x \, dx}{(x^2+a^2)(x^2+b^2)} \quad (a > b > 0) \quad \text{and} \quad \int_0^{\infty} \frac{\cos ax \, dx}{x^2+1} \quad (a > 0)$$