

# Exam 3

MAC 2313—CALCULUS III, SPRING 2017

(NEATLY!) PRINT NAME: \_\_\_\_\_

## Read all of what follows carefully before starting!

- This test has **6 problems** (12 parts total) and is worth **100 points**. *Please be sure you have all the questions before beginning!*
  - The exam is closed-note and closed-book. You may **not** consult with other students, and **no** calculators may be used!
  - Show all work clearly in order to receive full credit. Points will be deducted for incorrect work, and unless otherwise stated, no credit will be given for a correct answer without supporting calculations. **No work = no credit!** (unless otherwise stated)
  - You may use appropriate results from class and/or from the textbook as long as you fully and correctly state the result and where it came from.
    - If you use a result/theorem, you have to state *which* result you're using and explain *why* you're able to use it!
  - You **do not** need to simplify results, unless otherwise stated.
  - There is scratch paper at the end of the exam; you may also use the backs of pages or get more scratch paper from me.
  - Regions  $D$  in  $\mathbb{R}^2$  may be one and/or both of:
    - Type I if  $D = \{(x, y) : a \leq x \leq b, f_1(x) \leq y \leq f_2(x)\}$  for continuous functions  $f_1, f_2$
    - Type II if  $D = \{(x, y) : g_1(y) \leq x \leq g_2(y), c \leq y \leq d\}$  for continuous functions  $g_1, g_2$ .
  - Regions  $E$  in  $\mathbb{R}^3$  may be any and/or all of:
    - Type IB if  $E = \{(x, y, z) : (x, y) \in D, u_1(x, y) \leq z \leq u_2(x, y)\}$  for continuous functions  $u_1, u_2$
    - Type IIB if  $E = \{(x, y, z) : (y, z) \in D, u_1(y, z) \leq x \leq u_2(y, z)\}$  for continuous functions  $u_1, u_2$
    - Type III if  $E = \{(x, y, z) : (x, z) \in D, u_1(x, z) \leq y \leq u_2(x, z)\}$  for continuous functions  $u_1, u_2$ .
- Recall that " $\in$ " means "is an element of".

Question	1 (10 pts)	2 (25 pts)	3 (25 pts)	4 (20 pts)	5 (10 pts)	6 (10 pts)	Total (100 pts)
Points							

Do not write in these boxes! If you do, you get 0 points for those questions!

1. (5 pts ea.) Sketch each of the following regions and write each as the type designated. **There should be no integrals in your answers!**

(a)  $T_1$  = the triangle in  $\mathbb{R}^2$  with vertices  $(0, 0)$ ,  $(0, 2)$ ,  $(2, 2)$ ; as a Type I region.

(b)  $T_2$  = the same triangle as in (a); as a Type II region.

2. (a) (20 pts) Find the volume of the solid that lies under the paraboloid  $z = x^2 + y^2$  and above the region  $D$  in the  $xy$ -plane bounded by the line  $y = 2x$  and the parabola  $y = x^2$ . **Simplify fully.**

SOLUTION:

Part (b) is on the next page

(b) (5 pts) Now, set up the integral you would use to get the volume of the solid in part (a) using *the other* order of integration. **Do not integrate!**


**Hint:** In (a), you found the solution by expressing  $D$  as *either* a Type I *or* a Type II region; here, your answer should be the integral which expresses  $D$  as *the other* type.

SOLUTION:

3. (a) (5 pts) Let  $D$  be a region in the  $xy$ -plane, let  $z = f(x, y)$  be the equation of a surface  $S$ , and let  $A(S)$  denote the surface area of the portion of  $S$  over the region  $D$ . Write the formula for  $A(S)$ .

$$A(S) = \underline{\hspace{10cm}}.$$

- (b) (5 pts) Express the below region in polar coordinates.



SOLUTION:


Part (c) is on the next page

- (c) (15 pts) Find the area of the part of the paraboloid  $z = x^2 + y^2$  that lies under the plane  $z = 9$  and above the plane  $z = 1$ . **Hint:** This is part three of a problem; perhaps the other two parts help?

SOLUTION:

4. (20 pts) Use a triple integral to find the volume of the tetrahedron  $T$  bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$  (see below).

SOLUTION:



5. (5 pts ea.) Rewrite each of the rectangular iterated integrals in the coordinate system specified. **Do not integrate!**

(a)  $\int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \int_{x^2+y^2}^{2-x^2-y^2} \sqrt{x^2+y^2} dz dy dx$ ; in cylindrical coordinates.

SOLUTION:

Part (b) is on the next page



(b)  $\int_{-3}^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} \int_0^{\sqrt{9-x^2-y^2}} z \sqrt{x^2 + y^2 + z^2} dz dy dx$ ; in spherical coordinates.

SOLUTION:

6. (a) (5 pts) Compute the Jacobian of the transformation from rectangular coordinates to cylindrical coordinates in  $\mathbb{R}^3$ . **Simplify fully.**

SOLUTION:

**Part (b) is on the next page**

- (b) Use the result from part (a) to derive the formula for the triple integral in cylindrical coordinates of a function  $f(x, y, z)$  over a region  $E$  in  $\mathbb{R}^3$ .

SOLUTION:

**Bonus 1:**

(a) (4 pts) Verify that

$$f(x, y) = \begin{cases} 4xy & \text{if } 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$


is a joint (probability) density function.

(b) (3 pts ea.) Find the expected values of  $X$  and  $Y$ .

SOLUTION:

**Bonus 2:** (2 pts ea.) In question 4, you used triple integration to find the volume of the tetrahedron  $T$  bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , and  $x + y + z = 1$  shown below. Rewrite this integral as an equivalent iterated integral in the other five orders. **Do not integrate!**

**Hint:** Upon completing this problem, you should have evaluated/written six integrals—one each for  $dx dy dz$ ,  $dx dz dy$ ,  $dy dx dz$ ,  $dy dz dx$ ,  $dz dx dy$ , and  $dz dy dx$ —and each should give the same volume for  $T$ .



SOLUTION:

## Scratch Paper